

Name: _____

Operative Dentistry D262

Bonus Quiz

June 13, 2011

Seat Number: _____

Number correct: _____

Graded by: _____

10 E

~~*CORRECTED*~~

The introduction of Dr. Iggy into RDII this year caught many students by surprise. Since I don't curve grades I decided to give the class an opportunity for grade improvement by giving a bonus quiz. When calculating the final quiz grade I will drop your lowest 2 quiz grades so taking this quiz can only help your final grade.

1. The words "overcarved" and "undercarved", "overcontoured" and "undercontoured" are used frequently by your laboratory instructors. Some of them are included in the criteria for evaluation of the restorations done in your exercises and practical examinations. If a surface is undercarved, the surface is left

A

- ☒ a. overcontoured b. undercontoured c. also undercarved d. none of the above

2. When constructing a full gold crown, future recession of the gingival tissue can be most effectively prevented by

- a. narrowing the occlusal table by one-sixth
☒ b. accurately reproducing the buccal and lingual tooth form
c. slightly overcontouring the tooth form in the gingival one-fifth
d. extending the margins of the crown 1 mm into the gingival sulcus

B

F

F

T

T

T

T

3. (True ☒ False) Dycal has a low pH.

4. (True ☒ False) In a Class V cavity preparation for amalgam, the flaring of the mesial and distal walls is an example of retention form.

5. (True ☒ False) Inadequate length of composite cure, hyperocclusion, and inadequate primer placement are all likely to cause post-operative tooth sensitivity when restoring Class I and Class II composite restorations.

6. (True ☒ False) The level to which a base is built should never compromise the desired cavity preparation depth resulting in inadequate restorative material thickness.

7. (True ☒ False) For best access when restoring two adjacent teeth, prepare the largest lesion first and fill the smaller first.

8. (True ☒ False) When treatment requires that both a Class II and Class V preparation must be accomplished on the same tooth, the operator is advised to first complete the Class II preparation and restoration before initiating the Class V.

9. Because the proximity of the coronoid process, access to the facial surface of maxillary molars, particularly second molars, is often limited. The operator can have the patient move the mandible to increase visibility. Which of the following mandibular movements will improve access to the facial surface of the maxillary second molar?

- a. Have the patient protrude the mandible into an edge to edge anterior tooth position
☒ b. Have the patient partially close and shift the mandible toward the tooth being operated on
c. Have the patient partially close and shift the mandible away from the tooth being operated on
d. Have the patient open as wide as possible

B

10. Dr. Iggy has come to realize that he may not have taken full advantage of his educational experience in dental school. He failed to make the connection between education (receiving a piece of paper he can frame and hang on the wall) and true learning which can result in life-long improvement. But he's determined to provide the best dentistry he can for his patients and has adopted a new philosophy for his practice. Mediocrity is not acceptable in his practice and when he finds himself getting lazy, cutting corners, blaming others and making excuses he reminds himself of something he heard early in his dental school years which he's modified to fit his practice.

OK is not an _____ ? _____ and definitely not good enough for my patients.

Fill in the blank with whatever you want and take Dr. Iggy's advice don't waste what's left of your learning opportunity.