Endodontic Lab, Quiz 4B

February 10, 2003

Name \	06	
Name		Seat No
Brancowing and		

In a clinical situation, every tooth treated for endodontics must be isolated with a rubber dam. List three reasons stated in the lab manual for the placement of the rubber dam on the tooth.

- 81. To couse a tree solly working held
- 82. Retroct the soft hissive
- 83. Protect the policent from swellbring retruments or

According to lab manual, the exact measurement of root canal(s) of a posterior tooth is the distances from

- 84. Cusp tip to 85. spicel foramer.
- 86. The clamp selection is very important when placing on the rubber dam. As stated in lab manual, which of the following clamps would not be suitable for incisors and cuspids. 0,00,9

A - No. 0 B - No. 9 C No. 14 D - No. 00

List at least 3 purposes stated in the lab manual why radiographic analysis of clinical endodontic procedures is performed throughout the endodontic treatment.

- 87. To Get exact measurements and no debits in not and
- 88. Hoke sive the Gutta Percha is placed in the end of the noit and.
- 89. Hoke some no wide one present ding

As stated in lab manual, list two consequences of not removing all the pulp tissue from the chamber when accessing a maxillary pre-molar in a clinical situation.

90. Discoloration
91. Pois

Questions 92 through 95 are True or False.

- 92. In endodontic therapy only the tooth to be treated is isolated with the rubber dam and it is also the tooth that receives the clamp (nue)
- 93. When taking radiographs during the endodontic procedure the rubber dam and the rubber dam frame are removed in order to obtain a good x-ray False
- 94. The maxillary and mandibular pre molars tilt 30° lingual to the long ax s. Tal
- 95. The lingual canal orifice of a two-rooted maxillary pre-molar is located just lingual to the central fossa. True

Pretreatment with a provisional restoration may be necessary to provide sufficient structure to successfully place the rubber dam clamp. List four means, stated in lab manual of providing this treatment.

- 96. Orthodontic Bands.
- 97. Temporary Restriction
- 98. Temporary crown
- 99. Amalgam Build-up.

What is another reason, in addition to the one in the Question above, the lab manual states that pretreatment of a tooth may be needed.

100. Make sure that the root Conal are already properly and the root will not fracture during treatment.